VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals



Journal https://frontlinejournal s.org/journals/index.ph p/fmspj

Copyright: Original content from this work may be used under the terms of the creative commons attributes 4.0 licence.



POSTOPERATIVE EFFECTIVENESS OF PREOPERATIVE PREPARATION AND TREATMENT OF PATIENTS AT THE TIME OF THE THIRD STAGE OF RETRO AND PELVIC PROLAPSE IN WOMEN

Submission Date: January 19, 2024, Accepted Date: January 24, 2024,

Published Date: January 29, 2024

Crossref doi: https://doi.org/10.37547/medical-fmspj-04-01-05

Matyakubov Bunyod Bakhromovich

Doctor Obstetrical Gynecologist, Tashkent Medical Academy Urgench Branch, Department Of Obstetrics And Gynecology, Uzbekistan

Kulmatov G'Anijon Otakhonovich

Student, Urgench Branch Of Tashkent Medical Academy, Urgench, Uzbekistan

Niyazmetov Rakhmatulla Ermatovich

Doctor Of Medicine, Professor, Department Of Obstetrics Gynecologists, Tashkent Institute Of Advanced Medical Studies, Uzbekistan

TYT AL YILLY

29

ABSTRACT

Genital prolapse remains one of the unsolved problems of modern gynecology.

The aim of the study is to improve the preoperative preparation and postoperative management of patients with grade III pelvic organ prolapse, using modern innovative technology in combination with drugs to reduce postoperative complications and relapses.

Materials and methods of research. The object of the study was 217 patients with grade III pelvic organ prolapse who underwent surgical treatment.

Results and discussion. 110 women with grade III PG received an improved method of preoperative preparation, including irrigation with cavitated low-frequency ultrasound with an aqueous 0.05% chlorhexidine solution in the amount of 200.0, up to 4 times, locally conjugated estrogen, and in the

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

presence of urinary tract infection, acute cystitis - once fosfomycin trimetabol 3.0 grams orally, in recurrent forms of phytopreparation (Canephron N) for 14.5 ±2.5 days.

Conclusion: The improved method of preoperative preparation of patients with PTO of the III degree contributed to the reduction of the days of preoperative preparation by 6.0±0.5 days, reduced interoperative blood loss by 2.0 times, reduced the duration of the operation by 37.0±6.35 minutes, the hospital stay decreased by an average of 3.5±0.5 days, and postoperative recurrences after 1 year decreased by 2.4 times and after 2 years by 2.8 times.

KEYWORDS

Pelvic organ prolapse, genital prolapse, irrigation, conjugated estrogen, relapse.

Introduction

Relevance of the problem: Genital prolapse (PH) is a pathological process characterized by the prolapse and prolapse of the internal genitalia to or beyond the vaginal opening. PH is both a medical and social problem, leading to a reduced quality of life (1,4,6,7,8,9,10,11,13).pathology creates discomfort, causes urinary incontinence, constipation, leads to a decrease in libido, i.e. problems associated with sexual dysfunctions arise. In more advanced cases of genital prolapse, Trophic ulcers of the mucous membrane of the anterior wall of the vagina and cervix appear. (3,12,14,16,18,19,20,21,23). One of the relevances of this pathology is due to its high prevalence, high index of infectious and

inflammatory diseases against the background of pronounced violation of the vaginal microbiocenosis and urinary tract infection, as well as a high frequency of recurrences in the (1,2,24,25,26,27).postoperative period According to the WHO, there has been a significant increase in the number of patients suffering from genital prolapse worldwide in recent years. Today, between 6.5% and 64.6% of women suffer from PH of varying severity at different stages of life and in postmenopause, and their number in the world could reach 63 million by 2030. According to the American Association of Urology (AUA), one in nine women in the general population needs surgical treatment of

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

prolapse and/or genital stress urinary incontinence. Every year, approximately 400,000 women in the United States in an economically developed country undergo surgery for PH, however, the incidence of long-term postoperative recurrences is up to 30%, and early postoperative purulent infectious complications are 4 to 9%, which requires a new search for a solution to this problem (9,28). The ongoing scientific study of optimization of preoperative preparation and postoperative management of patients with genital prolapse, the use of modern innovative technologies reduces interoperative, early, long-term complications and relapses in this pathology.

The aim of this study is to improve the preoperative preparation and postoperative management of patients with grade III pelvic organ prolapse by using modern innovative technology in combination with drugs to reduce postoperative complications and relapses.

Material and methods of research. The object of the study was 217 patients with grade III pelvic prolapse who underwent surgical organ treatment. The collection of materials, analysis and evaluation of the results of preoperative preparation and postoperative management of

patients with pelvic organ prolapse in a comparative aspect was also carried out in the period from 2019 to 2023 on the basis of the gynecological department of the Bukhara district maternity hospital, the regional perinatal center and the private clinic "Lorastom". General clinical, biochemical, instrumental and statistical research methods were used in the study.

Discussion of the results. To assess the effectiveness of preoperative preparation and postoperative management of patients with grade III pelvic organ prolapse retro and prospective group, the object of the study was 217 women with grade III prolapse. All 217 patients with grade III pelvic organ prolapse were divided into two groups depending on the method of preoperative preparation and postoperative management. Group I A (1A) - (retrospectivecontrol) - 107 women with grade III PG, who received traditional preoperative preparation: vaginal debridement with a 1% alcohol solution of chlorophyllipt, ointment tamponade with Levomycol and Ovestin (micronized estriol 0.5 mg, vaginal suppositories 1 suppository per night) for 10±1.5 days. In the presence of urinary tract infection, recurrent and acute cystitis, the antibiotic Ceftriaxone was received 1.0x2 times

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

for 7.0±2 days. Rehabilitation of gynecological and somatic diseases.

The first B group (1B) - (prospective) - 110 women, also with stage III PH, who received an improved method of preoperative preparation and postoperative treatment. In the preoperative preparation, the FOTEC AK 101 device was used with cavitated low-frequency ultrasound with an aqueous 0.05% chlorhexidine solution in the amount of 200.0, up to 4 times, the level of ultrasonic vibrations was 50 units and the duration of exposure was 3 minutes. After 8 hours, 1.0 g of conjugated estrogen cream (femistron) was smeared on the entire prolapsed surface of the vaginal and uterine mucosa once a day, for 4.0±0.5 days. In the postoperative period, a three-component drug was received (Metostyl, 1 in the vagina at night for 5 days). In the presence of urinary tract infection, acute cystitis, a single dose of phosfomycin trimetabol 3.0 grams orally was given, in recurrent forms of phyto preparation (Canephron N) for 14.5 ±2.5 days. Studying the causes of PTO of the third degree in the retrospective and prospective groups from a comparative perspective, the following factors played a significant role: the number of pregnancies and births, concomitant somatic

diseases, estrogen deficiency in the menopausal period, profession and age. The peak incidence of PTO grade III occurred at the age of 40 to 59 years. At this age, this pathology occurred in the retrospective group in 62 (57.9%), in the prospective group in 74 (67.2%). It should be noted that in patients with PTO of the third degree of both groups, this pathology was mainly observed after the third birth. In the retrospective group after III-IV births, PTO grade III was detected in 72 (67.2%), and in the prospective group in 81 (73.6%). 88 (82.2%) in the retrospective group and 89 (80.9%) in the prospective group gave birth through the vaginal canal, which is one of the main risk factors for the development of PT. As indicated in the literature review, the influence of this factor is reflected in studies aimed at studying the so-called "bloating" phenomenon: 3D ultrasound after childbirth with the Valsalva test, in which a "swelling" of the muscle suspending the anus is detected (4,6).

In the menopausal period, grade III PTO was observed in 63 (58.8%) in the retrospective group and in 71 (64.5%) in the prospective group. During this period of life, there is not only a decrease in the systemic concentration of estrogens and their receptors for estrogens,

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

which are located in the nuclei of connective tissue cells, all this leads to a change in the properties of collagen. The observed estrogen deficiency in tissue and serum depends not only on the concentration of the hormone, but also on the level of expression and the receptor group. In the preoperative preparation, we paid special attention to the prescription of conjugated estrogens.

In 31 (28.9%) patients in the retrospective group and in 24 (21.8%) in the prospective group with stage III PTO, chronic cough was observed, i.e. chronic pathology of the bronchopulmonary system, and, accordingly, in 15 (14.0%) and in 11 (10%) complications after a coronavirus infection, which caused frequent dry cough. All these conditions lead to an increase in intraabdominal pressure and deterioration of the functional capacity of the pelvic tissue, which contributes not only to the development of PTO, but also to the deterioration of its course. The common somatic pathology most and complication in patients with stage III PTO was UTI, i.e. cystitis and pyelonephritis, which occurred in 42 (39.2%) cases in the retrospective group and in 48 (43.6%) cases in the prospective group, as well as cardiovascular diseases, such as

stage I-II hypertension, in 21 (19.6%) in the retrospective group and in 19 (17.2%) cases in prospective group. Vegetative-vascular the dystonia in 16 (14.9 %) and 12 (10.9%) cases, respectively <, and mild iron deficiency anemia in 62 (57.9%) in retrospective and in 71 (64.5%) p<0.05 cases in the prospective group, in a moderate-severe form, did not occur. The onset of PTO and pelvic organ prolapse does not occur immediately, this pathology develops slowly, patients pay attention when clinical symptoms appear, especially from the urinary tract. In the retrospective group, the first signs of clinical manifestation began after 3.5 years, i.e. the duration of PTO grade III ranged from 3.5 to 9 years with an average of 6.5±0.5 years p<0.05. And in the prospective group from 4 to 10 years, the average is 7.0±0.7 years, p<0.05. During this long time, due to the defect in the closure of the genital cleft (Fig. 1., Fig. 2., and Fig. 3), the obstetric trauma of the perineum or vagina is the main cause of the violation of the vaginal biocenosis. Changes in the biocenosis of the cervical canal are accompanied by inflammation, vaginosis and a decrease in local immunity with the subsequent formation of decubital ulcers (Fig. 2), and all this requires thorough preoperative preparation. In 34 (31.7%) patients with stage III

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

PTO in the prospective group, it was complicated by trophic ulcers of the mucous membrane of the anterior wall of the vagina and cervix. Before the start of debridement, the size of the ulcers varied from 2.0 to 6 cm2 and was characterized by pale vitreous granulations and flaccid epithelialization, purulent discharge from the

ulcer, and infected margins. The chronicity of vaginal inflammatory processes, the development of dysplasia and recurrent changes in the biocenosis have an extremely negative effect on the technique of the operation, the healing process increases the frequency of early and distant postoperative recurrences.







Rice. 1 cervical prolapse after amputation.

Rice. 2 initial stage of "decubital ulcers

Rice. 3 cervical polyps of the uterus

As described above, the vagina was treated with cavitated low-frequency ultrasound with an chlorhexidine solution in aqueous 0.05% advanced preoperative preparation, 2 sessions for grade II vaginal cleanliness, 3 sessions for grade III–IV, and 4-6 sessions for decubital ulcers. After conducting scientific research and studying the vaginal biotope, we fully confirm the data obtained in 2019 "Human Microbiome Project" of the vaginal biotope that there are 5 main classes

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727












Publisher: Frontline Journals

of bacterial communities in the vagina and of them 4 lactobacilli (L. iners, L. crispatus, L. gasseri or L. jensenii) and another 1 obligate anaerobes and they determine the cleanliness of the vagina. After two and three sessions, the state of the vaginal microbiocenosis improved, with repeated analyses the proportion of lactobacillus L. Crispatus increased from 53% to 78%; L. iners with 62% to 87% of cases, and L. gasseri with 51% to 74% of cases, L. jensenii from 43% to 65%. Changes in the vaginal biotope after cavitated irrigation with chlorhexidine in patients with grade III PH.

Diagram 1



After four sessions, the size of the decubital ulcers decreased by 2.1 times, from 1.0 to 3 cm2, which created good conditions for the operation. It should be noted that in only one patient, the size of decubital ulcers decreased to 10%, as she

suffered from a latent form of diabetes mellitus. The use of conjugated estrogen (femistron) trophism, improved tissue improved technique of surgery, reduced interoperative blood loss by 2.0 times p<0.05, reduced injuries

Volume 04 Issue 01-2024

35

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

and reduced the time of surgical intervention by 37.0 ±6.5 min. Femistron contributes to the restoration of normal vaginal epithelium in its atrophic changes in peri- and postmenopausal periods, normalization of vaginal microflora, thereby increasing the resistance of the vaginal epithelium to infectious and inflammatory processes. Against the background of an improvement in the outcome of the operation, the time spent in the hospital was reduced from an average of 7 days in the retrospective group to 4 days in the prospective group, i.e. by 1.7 times.

In the prospective group of 48 (43.6%) patients in preoperative preparation for acute cystitis, 9 (8.1%) received a single dose of fosfomycin trimetabol, with chronic and recurrent forms of BMI, 39 (35.4%) received the herbal drug Canephron N. This drug contains components - centaury, lovage and rosemary. It is all-natural and has an evidence base in terms of efficacy. All 39 patients were prescribed 2 tablets 3 times for 10 days before and after surgery. As a result, none of the 39 patients experienced a recurrence of UTIs in the early postoperative

period and during the first 3 months. In addition, intestinal dysbiosis decreased by 15 times compared to the retrospective group, where antibiotic therapy was received, and amounted to 2 (1.8%) cases in the prospective group. Thus, the prescription of the herbal drug Canephron N is effective and safe.

After preoperative preparation, all 110 patients of the prospective group underwent surgical treatment: anterior and posterior colporrhaphy with perineoleuvatoroplasty in 62 (56.3%); anterior and posterior colporrhaphy with perineoleuvatoroplasty and wind fixation in 36 (32.7%); anterior and posterior colporrhaphy with perineoleuvatoroplasty and cervical amputation 12 (10.9%) p<0.05.

The assessment of the effectiveness of preoperative preparation, surgical treatment and rehabilitation in a comparative version is shown in Table № 1.

Evaluation of the effectiveness of preoperative preparation, surgical treatment and rehabilitation in comparison with PG-III degree.

36

Table 1

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

	Patients with grade III PH	
С	Retrospective group (n=107)	Prospective group(n=110)
r i t e r i	Abs %	Abs %
Duration of the operation, min	105± 7.2 min*	68± 5.5 min*
Average blood loss ml	350± 60.0ml*	170± 35.0ml*
Increase in body temperature to 380	RONT	4 3,6*
Suture infiltrate after surgery		3 2,7*
Exacerbation of urinary tract infection aftersurgery	13	4 3,6*
Average postoperative bed day	7± 0.5 days	4± 0.5 days
Postoperative relapse after 1 year	12	5 4,5
Postoperative relapse after 2 year	19	7 6,3

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

Notes: Static confidence * p<0.05.

In the postoperative period, all patients received a three-component drug (Metostyl 1 suppository in the vagina at night for 5 days). Metostyl is an antibacterial, anti-inflammatory and antifungal drug in the form of vaginal suppositories. The use of metostil in the postoperative period after surgery for genital prolapse vaginal contributed to an increase in the rate of tissue regeneration and improved the effectiveness of surgical treatment of patients with prolapse of the internal genitalia. The use of metostil in the complex of preoperative preparation made it possible to reduce the duration of treatment by 1.7 times and the infiltration of sutures by 2.1 times. In the postoperative period, all patients received a three-component drug (Metostyl 1 vaginal suppositories at night for 5 days). Metostyl is an antibacterial, anti-inflammatory and antifungal drug in the form of vaginal suppositories. One suppository contains: 200 mg metronidazole, 160 mg clotrimazole, and 8 mg chlorhexidine. The use of mettostyle in the postoperative period after vaginal operations for grade III genital prolapse contributed to an increase in the rate of tissue regeneration and improves the effectiveness of surgical treatment

of patients with prolapse of the internal genital organs. The use of metostyle in the complex of preoperative preparation made it possible to reduce the duration of treatment by 1.7 times and the infiltration of sutures by 2.1 times compared to the retrospective group. Assessing the effectiveness of preoperative preparation and postoperative management of patients with retro and prospective pelvic organ prolapse of the third degree, we came to the following conclusions:

INDINGS

- The improved method of preoperative preparation of patients with PTO of the III degree contributed to the reduction of the days of preoperative preparation by 6.0±0.5 days, reduced interoperative blood loss by 2.0 times (by 180.0±32.0 ml), this method reduced the duration of the operation by 37.0±6.35 minutes, reduced the stay in the hospital by an average of 3.5±0.5 days, and reduced early postoperative purulent infectious complications by 2.1 times.
- 2) Postoperative recurrences decreased by 2.4 times after 1 year and by 2.8 times after 2 years.

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

and at the same time the quality of life improved in all patients.

3) The economic efficiency of preoperative preparation of the recommended method was 72.1% and postoperative – 40.6% p<0.05.

REFERENCES

- 1. Badal'yants D.A. Clinical features of genital prolapse in patients who need surgical treatment. 2020. S. 38-39.
- 2. Gafurov Zh.M. Improvement of sling plastic surgeries in women and assessment of their effectiveness. Abstract of the dissertation of Doctor (PhD) of Philosophy in Medical Sciences .Tashkent.2021.p.41
- **3.** Danilina O.A., Volkov V.G. Prevalence of pelvic among organ prolapse women reproductive age // Journal of new medical technologies, 2022 – vol. 29, No. 1 – p. 29-33.
- **4.** Orazov M.R., Silantyeva E.S., Soldatskaya R.A., Belkovskaya M.E. Efficacy of physiotherapy in the treatment of pelvic floor muscle insufficiency in women of reproductive age. Doctor.Ru. 2020; 19(8):71-76.
- 5. Radzinsky V.E., Orazov M.R., Tokar L.R. et al. Perineology. Aesthetic gynecology. Moscow, 2022-416p.

- **6.** Mattsson, N.K. [et al.] Pelvic organ prolapse surgery and quality of life-a nationwide cohort study // Am J Obstet Gynecol.- 2020.-Vol.222, №6.-P:588.
- 7. NICE Guidance Urinary incontinence and pelvic organ prolapse in women: management: NICE (2019)Urinary incontinence and pelvic organ prolapse in women: management. BJU Int. 2019 May; 123(5):777-803. doi: 10.1111/bju.14763. PMID: 31008559.
- Хамдамов И.Б. Клиническая оценка эффективности традиционного подхода лечения грыж передней брюшной стенки у женщин фертильного возраста // Вестник врача. -Самарканд 2022. № 2.2 (104).- С.65-70.
- 9. Хамдамов И.Б., Мирходжаев И.А. Хакимов М.Ш. Хамдамов Б.З. Эволюция использования полимерных имплантантов для герниопластики // Тиббиётда янги кун. – Ташкент; 2021,- №2 (34) C.-107-111.
- **10.**Khamdamov I.B., Khamdamov A.B. Differentiated approach to the choice of hernioplasty method in women of fertile age experimental (Clinical and study) //

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

- Тиббиётда янги кун. Бухоро, 2021.-№ 6 (38/1).-C. 112- 114.
- 11. Хамдамов И.Б., Хамдамов А.Б. Фертил ёшдаги аёлларда эндовидеохирургик герниопластика // Тиббиётда янги кун. Бухоро, 2021.-№6 (38/1) -С. 25-27.
- **12.** Хамдамов И.Б. Experimental determination of the extensibility of the anterior abdominal wall tissues at different times of pregnancy using various approaches to hernioplasty// Academicia: An International Multidisciplinary Research Journal Vol. 12, Issue 04, April 2022 SJIF 2022 = 8.252 P.193-201
- **13.** Хамдамов И.Б. Совершенствование тактических подходов в лечении грыж передней брюшной стенки у женщин фертильного возраста // Тиббиётда янги кун. Бухоро, 2022.-№10(48)- С. 338-342. 14. Хамдамов И.Б. Морфофункциональные особенности брюшного пресса у женщин репродуктивного возраста // Тиббиётда янги кун. Бухоро, 2022.-№3(41)- С. 223-227.
- **14.**Khamdamova M.T. Ultrasound features of three-dimensional echography in assessing the condition of the endometrium and uterine cavity in women of the first period of middle age using intrauterine contraceptives //

- Biology va tibbyot muammolari. Samarkand, 2020. - No. 2 (118). - P.127-131.
- **15.** Khamdamova M. T. Ultrasound assessment of changes in the endometrium of the uterus in women of the first and second period of middle age when using intrauterine and oral contraceptives // Биомедицина ва амалиёт журнали. - Ташкент, 2020. - №2. - 8 часть. -C.79-85.
- **16.**Khamdamova M. Т. Anthropometric characteristics of the physical status of women in the first and second period of middle age // A new day in medicine. Tashkent, 2020. - № 1 (29). - C.98-100.
- 17. Khamdamova M.T. Age-related and individual variability of the shape and size of the uterus according to morphological and ultrasound studies // News of dermatovenereology and reproductive health. - Tashkent, 2020. - No. 1-2 (88-80). - P.49-52.
- **18.**Khamdamova M. T. Anthropometric characteristics of the physical status of women in the first and second period of middle age // Тиббиётда янги Ташкент, 2020. - № 1 (29). - С.98-100.
- 19. Хамдамова M.T. Возрастная И индивидуальная изменчивость формы и размеров матки по данным

40

VOLUME 04 ISSUE 01 Pages: 29-41

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











Publisher: Frontline Journals

- морфологического ультразвукового Новости исследований дерматовенерологии и репродуктивного здоровья. - Ташкент, 2020. - № 1-2 (88-80). - C.49-52.
- **20.** Хамдамова M.T. Ультразвуковая особенности трехмерный эхографии в оценке состояния эндометрия и полости матки у женщин первого периода среднего возраста применяющие внутриматочные контрацептивные средства // Биология ва тиббиёт муаммолари. - Самарканд, 2020. -Nº2 (118). - C.127-131.
- 21. Khamdamova M. T. Ultrasound assessment of changes in the endometrium of the uterus in women of the first and second period of middle age when using intrauterine and oral contraceptives // Биомедицина ва амалиёт журнали. - Ташкент, 2020. - №2. - 8 часть. -C.79-85.
- 22. Хамдамова M.T. Особенности ультразвуковых параметров матки у женщин первого и второго периода среднего возраста применяющие

- контрацептивные инъекционные средства // Тиббиётда янги кун. - Ташкент, 2020. - № 2/1 (29/1). - C.154-156.
- **23.** Хамдамова M.T. Особенности ультразвукового изображения матки и яичников у женщин второго периода среднего возраста применяющие комбинированные оральные контрацептивные средства // Тиббиётда янги кун. - Ташкент, 2020. - № 2 (30). - С. 258-261.
- **24.** Хамдамова M.T. Индивидуальная изменчивость матки и яичников у женщин применяющие и не использующие различные контрацептивные виды средства // Тиббиётда янги кун. - Ташкент, 2020. - № 3 (31). - C. 519-526.
- **25.** Khamdamova M. T. Echographic features variability in the size and shape of the uterus and ovaries in women of the second period of adulthood using various contraceptives // Asian Journal of Multidimensional Research -2020. - N9 (5). - P.259-263.

Volume 04 Issue 01-2024

41