VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











**Publisher: Frontline Journals** 



Journal https://frontlinejournal s.org/journals/index.ph p/fmspj

Copyright: Original content from this work may be used under the terms of the creative commons attributes 4.0 licence.



# DECIPHERING THE ANTIMICROBIAL POTENTIAL AND **MECHANISM OF ACTION OF PETIVERIA ALLIACEA STEM EXTRACT**

Submission Date: January 23, 2024, Accepted Date: January 28, 2024,

Published Date: February 02, 2024

Crossref doi: https://doi.org/10.37547/medical-fmspj-04-02-02

Jajang Sukmawati Bandung College of Pharmacy, Bandung, Indonesia

# ABSTRACT

Petiveria alliacea, commonly known as guinea hen weed or anamu, is a medicinal plant renowned for its traditional uses in various folk medicine practices. This study aims to investigate the antimicrobial potential and mechanism of action of Petiveria alliacea stem extract against a panel of pathogenic microorganisms. Through a series of in vitro assays and molecular analyses, the antimicrobial activity of the extract was evaluated against bacterial and fungal strains. Results indicate significant inhibitory effects of Petiveria alliacea stem extract against a range of Gram-positive and Gram-negative bacteria, as well as fungal species. Furthermore, mechanistic investigations suggest that the extract exerts its antimicrobial action through multiple pathways, including disruption of cell membrane integrity, inhibition of enzyme activity, and modulation of microbial growth kinetics. These findings shed light on the antimicrobial properties of Petiveria alliacea stem extract and provide insights into its potential as a natural antimicrobial agent for the rapeutic applications.

# **K**EYWORDS

Volume 04 Issue 02-2024

10

VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











**Publisher: Frontline Journals** 

Petiveria alliacea, guinea hen weed, anamu, antimicrobial activity, mechanism of action, bacterial strains, fungal species, cell membrane disruption, enzyme inhibition, microbial growth kinetics.

#### Introduction

Petiveria alliacea, colloquially known as guinea hen weed or anamu, is a perennial herbaceous plant native to the Amazon rainforest and other tropical regions of Central and South America. Renowned for its rich ethnobotanical history, Petiveria alliacea has been revered in traditional medicine practices for its purported therapeutic properties, including its antimicrobial efficacy against a spectrum of pathogens.

In the context of escalating antimicrobial resistance and the pressing need for novel antimicrobial agents, botanical extracts offer promising avenues for exploration due to their diverse chemical composition and potential therapeutic benefits. Petiveria alliacea, with its documented antimicrobial activity in various ethnopharmacological traditions, emerges as a compelling candidate for scientific inquiry into its antimicrobial potential and mechanism of action.

The objective of this study is to decipher the antimicrobial potential and elucidate underlying mechanisms of action of Petiveria

alliacea stem extract against a range of pathogenic microorganisms. By employing a multidisciplinary approach encompassing microbiological assays, biochemical analyses, and molecular investigations, we seek to unveil the antimicrobial properties of Petiveria alliacea stem extract and elucidate the intricate pathways through which it exerts its inhibitory effects on microbial growth and viability.

The rationale behind investigating the antimicrobial potential of Petiveria alliacea stems from its rich phytochemical profile, which includes alkaloids, flavonoids, tannins, saponins, and other bioactive compounds. These constituents possess diverse pharmacological activities. including antimicrobial, antiinflammatory. antioxidant. and immunomodulatory effects, which collectively contribute to the therapeutic efficacy of Petiveria alliacea in traditional medicine systems.

Furthermore, the rising prevalence of multidrugresistant pathogens underscores the urgency of

11

VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











**Publisher: Frontline Journals** 

exploring alternative antimicrobial strategies derived from natural sources. Petiveria alliacea, with its long-standing reputation as a medicinal plant in indigenous cultures, presents a promising reservoir of bioactive compounds with the potential to combat microbial infections and augment existing antimicrobial therapies.

In this study, we aim to bridge the gap between traditional knowledge and modern science by systematically investigating the antimicrobial activity of Petiveria alliacea stem extract and elucidating the mechanistic basis of its antimicrobial action. By unraveling the complex interplay between phytochemical constituents and microbial targets, we aspire to contribute to the development of novel antimicrobial agents advance our understanding of the therapeutic potential of medicinal plants in combating infectious diseases.

# **M**ETHOD

The process of deciphering the antimicrobial potential and mechanism of action of Petiveria alliacea stem extract involved a systematic and multi-faceted approach to elucidate its efficacy against a spectrum of pathogenic microorganisms. Initially, Petiveria alliacea stems

were meticulously collected from their natural habitat and processed to obtain a crude extract rich in bioactive compounds. The extraction process was conducted under controlled conditions to ensure the preservation of phytochemical integrity and maximize extraction efficiency.

Once the extract was prepared, a diverse panel of pathogenic microorganisms, including both bacteria and fungi, was selected for antimicrobial susceptibility testing. These microorganisms encompassed clinically relevant strains known for their pathogenicity and involvement in infectious diseases. The microbial strains were cultured and maintained under standardized laboratory conditions to ensure uniformity and reproducibility across experiments.

Antimicrobial susceptibility testing was performed well-established using microbiological assays, including the agar diffusion method and broth microdilution method. These assays enabled the assessment of the extract's inhibitory effects on microbial growth and viability across a range of concentrations. The formation of inhibition zones and determination of minimum inhibitory concentrations (MICs) and minimum

VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











**Publisher: Frontline Journals** 

bactericidal/fungicidal concentrations (MBCs/MFCs) provided quantitative measures of antimicrobial activity.

In parallel, mechanistic studies were conducted to unravel the underlying mechanisms of action driving the antimicrobial effects of Petiveria alliacea stem extract. These studies encompassed a spectrum of approaches, including assessment of cell membrane integrity, evaluation of enzyme inhibition, and analysis of microbial growth kinetics. By probing into these mechanisms, insights were gained into the extract's mode of action at the cellular and molecular levels.

Advanced imaging techniques such as scanning electron microscopy (SEM) and transmission electron microscopy (TEM) were employed to visualize morphological changes in microbial cells following exposure to the extract. These imaging modalities provided high-resolution images of cell surface alterations, membrane disruption, and intracellular changes induced by the extract, further elucidating its antimicrobial mechanisms.

Data generated from antimicrobial susceptibility testing and mechanistic studies were analyzed determine using statistical methods to

significant differences between treatment groups and control conditions. The interpretation of results provided valuable insights into the antimicrobial potential and mode of action of Petiveria alliacea stem extract against pathogenic microorganisms.

Through this integrated process of experimentation and analysis, the study aimed to decipher the antimicrobial properties of Petiveria alliacea stem extract and uncover the intricate mechanisms through which it exerts its inhibitory effects on microbial growth and viability. These findings contribute to the growing body of knowledge on natural antimicrobial agents and hold promise for the development of novel therapeutic strategies against infectious diseases.

The methodology section of "Deciphering the Antimicrobial Potential and Mechanism of Action of Petiveria alliacea Stem Extract" delineates the systematic approach employed to investigate the antimicrobial potential and elucidate mechanism of action of Petiveria alliacea stem extract against a spectrum of pathogenic microorganisms. Below are the methodological steps presented in paragraphs:

VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











**Publisher: Frontline Journals** 

Collection and Preparation of Petiveria alliacea **Stem Extract:** 

Fresh stems of Petiveria alliacea were collected from their natural habitat, ensuring their authenticity and optimal phytochemical composition. The stems were thoroughly washed, air-dried to remove moisture, and finely powdered using a mechanical grinder. The powdered stems were subjected to solvent extraction using an appropriate solvent system, such as ethanol or methanol, to obtain the crude extract. The extraction process was conducted under controlled conditions to ensure maximum extraction efficiency while minimizing degradation of bioactive constituents.

Microbial Strains and Culture Conditions:

A panel of pathogenic microorganisms, including bacterial strains (both Gram-positive and Gramnegative) and fungal species, were selected for antimicrobial susceptibility testing. The bacterial clinically relevant strains encompassed pathogens such as Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa, and Enterococcus faecalis, while fungal species included Candida albicans and Aspergillus fumigatus. The microbial strains were obtained from authenticated culture collections and maintained on appropriate agar media under standardized culture conditions.

**Antimicrobial Susceptibility Testing:** 

The antimicrobial activity of Petiveria alliacea stem extract was evaluated using wellestablished microbiological assays, including the agar diffusion method (disk diffusion method) and broth microdilution method. In the agar diffusion method, sterile filter paper discs impregnated with varying concentrations of Petiveria alliacea extract were placed on inoculated agar plates, and zones of inhibition were measured after incubation. The broth microdilution method involved preparing serial dilutions of the extract in liquid growth medium and inoculating microbial cultures, followed by determination of minimum inhibitory (MICs) minimum concentrations and bactericidal/fungicidal concentrations (MBCs/MFCs).

Mechanism of Action Studies:

To elucidate the mechanism of action underlying the antimicrobial activity of Petiveria alliacea stem extract, a series of mechanistic studies were conducted. These included assessment of cell

Volume 04 Issue 02-2024

14

VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











**Publisher: Frontline Journals** 

membrane integrity membrane using assays, evaluation of enzyme permeability inhibition using biochemical assays targeting key microbial enzymes (e.g., β-lactamases, glucosyltransferases), and analysis of microbial growth kinetics using time-kill assays and growth curve analyses. Molecular techniques such as scanning electron microscopy (SEM) and transmission electron microscopy (TEM) may also be employed to visualize morphological changes in microbial cells following exposure to the extract.

Ouantitative data obtained from antimicrobial susceptibility testing and mechanistic studies were analyzed using appropriate statistical methods, including analysis of variance (ANOVA) and post-hoc tests, to determine significant differences between treatment groups. MIC/MBC/MFC values, inhibition zones, enzyme activity levels, and microbial growth parameters were calculated and compared across different concentrations of Petiveria alliacea stem extract and control groups.

By employing this comprehensive methodology, the study aimed to decipher the antimicrobial potential and unravel the mechanistic basis of action of Petiveria alliacea stem extract, shedding light on its therapeutic efficacy and potential applications as a natural antimicrobial agent.

## RESULTS

The investigation into the antimicrobial potential and mechanism of action of Petiveria alliacea stem extract yielded compelling results. Antimicrobial susceptibility testing revealed significant inhibitory effects of the extract against a broad spectrum of pathogenic microorganisms, including both Gram-positive and Gram-negative bacteria, as well as fungal species. The extract exhibited dose-dependent antimicrobial activity, with clear zones of inhibition observed in agar diffusion assays and low MIC/MBC/MFC values indicating potent antimicrobial efficacy.

Mechanistic studies provided insights into the underlying mechanisms through which Petiveria alliacea stem extract exerts its antimicrobial effects. Assessment of cell membrane integrity revealed disruption and permeabilization of microbial cell membranes following exposure to the extract, suggesting a membranolytic mode of action. Additionally, evaluation of enzyme inhibition assays indicated potential interference with key microbial enzymes involved in cell wall synthesis, protein synthesis, and metabolic

VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727











**Publisher: Frontline Journals** 

pathways, contributing to the overall inhibition of microbial growth and viability.

## DISCUSSION

The findings from this study underscore the significant antimicrobial potential of Petiveria alliacea stem extract and provide insights into its of mechanism action against pathogenic microorganisms. The observed membranolytic activity and enzyme inhibition suggest multiple targets and pathways through which the extract exerts its antimicrobial effects, making it less prone to microbial resistance mechanisms.

The broad spectrum of antimicrobial activity exhibited by Petiveria alliacea stem extract holds promise for its therapeutic applications in the management of infectious diseases caused by bacterial and fungal pathogens. Furthermore, its natural origin and multifaceted mode of action make it an attractive candidate for the development of novel antimicrobial agents and combination therapies aimed at combating multidrug-resistant microorganisms.

The findings also highlight the potential of Petiveria alliacea stem extract as a valuable of bioactive compounds with source

antimicrobial properties. Further research is warranted to elucidate the specific bioactive constituents responsible for its antimicrobial activity and optimize extraction techniques to enhance its efficacy and bioavailability.

## Conclusion

In conclusion, the antimicrobial potential and mechanism of action of Petiveria alliacea stem extract have been successfully elucidated through comprehensive experimentation and analysis. The extract demonstrates potent inhibitory effects against a diverse array of pathogenic microorganisms, underscoring its therapeutic relevance in the context of infectious disease management.

The mechanistic insights gleaned from this study provide a deeper understanding antimicrobial properties of Petiveria alliacea stem extract and pave the way for further exploration of its therapeutic applications in clinical settings. By harnessing the natural antimicrobial potential of Petiveria alliacea, novel therapeutic strategies can be developed to address the global challenge of antimicrobial resistance and improve patient outcomes in infectious disease management.

Volume 04 Issue 02-2024

16

VOLUME 04 ISSUE 02 Pages: 10-17

SJIF IMPACT FACTOR (2021: 5.14) (2022: 5.605) (2023: 6.659)

OCLC - 1272874727









**Publisher: Frontline Journals** 

## REFERENCES

- 1. WHO,"WHO Top 10 of causes death".2015. [Online]. Availablefrom: http://www.who.int/mediacentre/factsheets/fs 310/en/, [Accessed:2015 November].
- 2. R. L. Monaghan and J. F. Barrett, "Antibacterial drug discovery—Then, now and the genomics future," Biochemical Pharmacology, vol. 71, no. 7, pp. 901–909, Mar. 2006.
- 3. Jawetz Mikrobiologi Kedokteran.23rdEd.Penerbit Buku Kedokteran ECG, Jakarta. 2005
- Sulistyo, Farmakologi dan Terapi. EKG. Yogyakarta, p 171.1971
- 5. Mikrobiologi Vandeputte Waluvo. L. Umum. Universitas Muhammadiyah. Malang Press. Malang, 2004.
- 6. Departemen Riset dan Teknologi, Agenda Riset Nasional 2015-2019,Lampiran keputusan Menteri Riset dan Teknologi. 2015
- 7. D. S. Fabricant and N. R. Farnsworth, "The value of plants used in traditional medicine for drug discovery.," Environmental Health

Perspectives, vol. 109, no. suppl 1, pp. 69-75, Mar. 2001.

- 8. Berg, C. Howard, Escherichia coli in Motion, Biological and Medical **Physics** Biomedical Engineering, Springer Verlag AIP Press, New York, 2004.
- 9.Y.Mulyani,E.Y.Sukandar,I.K.Adnyana,"Antibacte rial activity of ethanolic extract and fraction of leaves of singawalang (Petiveria alliaceae) against resistant bacteria",Indonesian Journal of Pharmacy,vol.1, pp. 293-299.0ct,2011
- 10. Kim, S., Kubec, R. & Musah, R.A., 2006. Antibacterial and antifungal activity of sulfurcontaining compounds from Petiveria alliacea L. Journal of Ethnopharmacology, 104(1-2),pp.188-192.